

- [3] Distant S, Bjoro K, Hellum KB, Myrvang B, Berg JP, Skaug K, et al. Raised serum ferritin predicts non-response to interferon and ribavirin treatment in patients with chronic hepatitis C infection. *Liver* 2002;22:269–275.
- [4] Bonkovsky HL, Naishadham D, Lambrecht RW, Chung RT, Hoefs JC, Nash SR, et al. Roles of iron and HFE mutations on severity and response to therapy during retreatment of advanced chronic hepatitis C. *Gastroenterology* 2006;131:1440–1451.
- [5] Lebray P, Zylberberg H, Hue S, Poulet B, Carnot F, Martin S, et al. Influence of HFE gene polymorphism on the progression and treatment of chronic hepatitis C. *J Viral Hepat* 2004;11:175–182.
- [6] Sebastiani G, Walker AP. HFE gene in primary and secondary hepatic iron overload. *World J Gastroenterol* 2007;13:4673–4689.

Carine Fillebeen
Lady Davis Institute for Medical Research

Sir Mortimer B. Davis Jewish General Hospital,
3755 Cote-Ste-Catherine Road, Montreal,
Quebec, Canada H3T 1E2

Kostas Pantopoulos*
Lady Davis Institute for Medical Research
Sir Mortimer B. Davis Jewish General Hospital
3755 Cote-Ste-Catherine Road, Montreal
Quebec, Canada H3T 1E2

Department of Medicine, McGill University, Canada
* Tel.: +1 (514) 340 8260x5293; fax: +1 (514) 340 7502
E-mail address: kostas.pantopoulos@mcgill.ca

Hepatocyte iron accumulation: A new string to ribavirin's antiviral bow?

To the Editor:

Building on previous work, the recent article by Fillebeen and Pantopoulos elegantly demonstrates a role for iron as an inhibitor of hepatitis C virus (HCV) replication [1]. Using HCV-infected Huh7.5.1 hepatoma cells, the authors show a dose-dependent reduction in the expression of HCV viral proteins and RNA upon exogenous administration of iron. Moreover, the anti-viral effect of iron was attributed to the direct inhibition of the HCV RNA polymerase NS5B by iron [2].

However, the findings outlined in this article appear contrary to the prevailing perception of the role of iron in HCV infection. Disordered iron homeostasis is a frequent finding in HCV patients, and may be associated with adverse clinical outcomes [3]. An increased propensity to hepatic decompensation, an increased incidence of hepatocellular carcinoma, and a reduced response to treatment have all been reported in association with excess serum or hepatic iron [4–6]. The authors acknowledge these issues, and conclude by doubting whether iron-mediated inhibition of HCV replication would represent a realistic therapeutic target. These sentiments were echoed in the accompanying editorial [7].

The nucleoside analogue ribavirin forms a key component of the current standard of care for HCV treatment, in combination with pegylated interferon alpha. Despite its importance in augmenting treatment response and preventing relapse, its antiviral mechanism of action remains elusive [8]. Curiously, significant hepatic iron accumulation has been reported in HCV patients receiving ribavirin monotherapy for greater than 6 months [9,10]. These findings were attributed to the well-documented, dose-dependent haemolysis caused by ribavirin. However, excess iron accumulated predominantly in hepatocytes, rather than in phagocytic Kupffer cells, as might be expected following intravascular haemolysis [10]. Although hepatic iron accumulation during treatment was not associated with changes in liver transaminases, it is unclear whether treatment response was altered [10].

Iron accumulation in chronic HCV infection appears harmful. Given the findings by Fillebeen et al., it would be difficult to dis-

count a potential role for iron in hepatocyte HCV eradication, which may be transiently facilitated by ribavirin therapy. At the very least, this potential mechanism of ribavirin action merits further investigation.

Financial support

GI Research fund, Mater Hospital.

Conflict of interest

The authors who have taken part in this study declared that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

References

- [1] Fillebeen C, Pantopoulos K. Iron inhibits replication of infectious hepatitis C virus in permissive Huh7.5.1 cells. *J Hepatol* 2010;53:995–999.
- [2] Fillebeen C, Rivas-Estilla AM, Bisaillon M, Ponka P, Muckenthaler M, Hentze MW, et al. Iron inactivates the RNA polymerase NS5B and suppresses subgenomic replication of hepatitis C virus. *J Biol Chem* 2005;280:9049–9057.
- [3] Piperno A, Vergani A, Malosio I, Parma L, Fossati L, Ricci A, et al. Hepatic iron overload in patients with chronic viral hepatitis: role of HFE gene mutations. *Hepatology* 1998;28:1105–1109.
- [4] Fujita N, Sugimoto R, Urawa N, Araki J, Mifuji R, Yamamoto M, et al. Hepatic iron accumulation is associated with disease progression and resistance to interferon/ribavirin combination therapy in chronic hepatitis C. *J Gastroenterol Hepatol* 2007;22:1886–1893.
- [5] Bassett SE, Bisceglie AMD, Bacon BR, Sharp RM, Govindarajan S, Hubbard GB, et al. Effects of iron loading on pathogenicity in hepatitis C virus-infected chimpanzees. *Hepatology* 1999;29:1884–1892.
- [6] Chapoutot C, Esslimani M, Joomaye Z, Ramos J, Perney P, Laurent C, et al. Liver iron excess in patients with hepatocellular carcinoma developed on viral C cirrhosis. *Gut* 2000;46:711–714.
- [7] Mueller S. Increased iron in HCV infection: collateral damage or antiviral defense? *J Hepatol* 2010;53:990–992.
- [8] Feld JJ, Hoofnagle JH. Mechanism of action of interferon and ribavirin in treatment of hepatitis C. *Nature* 2005;436:967–972.

Letters to the Editor

- [9] Di Bisceglie AM, Bacon BR, Kleiner DE, Hoofnagle JH. Increase in hepatic iron stores following prolonged therapy with ribavirin in patients with chronic hepatitis C. *J Hepatol* 1994;21:1109–1112.
- [10] Fiel MI, Schiano TD, Guido M, Thung SN, Lindsay KL, Davis GL, et al. Increased hepatic iron deposition resulting from treatment of chronic hepatitis C with ribavirin. *Am J Clin Pathol* 2000;113:35–39.

John D. Ryan*

Centre for Liver Disease, Mater Misericordiae University Hospital
Dublin, Ireland
School of Medicine and Medical Science, University College
Dublin, Ireland

* Tel.: +353 861701667, fax: +353 18034058

E-mail addresses: jdpryan@yahoo.co.uk, johnryan@physicians.ie

John Crowe

Centre for Liver Disease, Mater Misericordiae University Hospital
Dublin, Ireland
School of Medicine and Medical Science, University College
Dublin, Ireland

Reply to: Hepatocyte iron accumulation: A new string to ribavirin's antiviral bow?

This is a reply to the letter to the Editor by Ryan and Crowe:

We thank Drs. J.D. Ryan and J. Crowe for their comments to our recent article [1]. In their letter, they discuss earlier reports showing that prolonged treatment of HCV-infected patients with ribavirin associates with hepatic iron deposition, mainly in hepatocytes [2,3]. This is largely attributed to *extravascular* hemolysis, a known side effect of ribavirin ([2] and references therein) even though one would expect a predominant distribution of excess iron within Kupffer cells. The authors further hypothesise that accumulation of iron within hepatocytes may potentiate the antiviral efficacy of this therapeutic regimen, considering that iron inhibits the replication of HCV in permissive Huh7.5.1 cells [1].

We completely agree that this hypothesis merits further exploration. It will be interesting to examine the effects of ribavirin in the infectious HCV replicon model, with or without exogenous iron administration. Nevertheless, given the complexity of the pharmacological responses to ribavirin *in vivo*, which impinges on systemic iron traffic, we should be cautious in extrapolating data from this cell culture model. Moreover, the iron deposition in hepatocytes of patients did not appear to affect the “biochemical or histologic response to ribavirin therapy” over a period of 9 months, even though sustained virological response (SVR) was not assessed [2]. A more recent prospective study evaluated how iron perturbations affect combination therapy with pegylated interferon and ribavirin [4]. The data revealed a negative correlation between iron load in Kupffer cells and the prospect of achieving SVR, and did not provide any evidence for potential benefits of hepatocyte iron accumulation. Further clinical studies with HCV-infected patients and, possibly, also experiments with humanized mouse models for HCV infection [5], may provide a more clear view on how hepatocyte iron accumulation interferes with antiviral therapy.

Conflict of interest

The authors who have taken part in this study declared that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

References

- [1] Fillebeen C, Pantopoulos K. Iron inhibits replication of infectious hepatitis C virus in permissive Huh7.5.1 cells. *J Hepatol* 2010;53:995–999.
- [2] Fiel MI, Schiano TD, Guido M, Thung SN, Lindsay KL, Davis GL, et al. Increased hepatic iron deposition resulting from treatment of chronic hepatitis C with ribavirin. *Am J Clin Pathol* 2000;113:35–39.
- [3] Di Bisceglie AM, Bacon BR, Kleiner DE, Hoofnagle JH. Increase in hepatic iron stores following prolonged therapy with ribavirin in patients with chronic hepatitis C. *J Hepatol* 1994;21:1109–1112.
- [4] Ferrara F, Ventura P, Vegetti A, Guido M, Abbati G, Corradini E, et al. Serum ferritin as a predictor of treatment outcome in patients with chronic hepatitis C. *Am J Gastroenterol* 2009;104:605–616.
- [5] Washburn ML, Bility MT, Zhang L, Kovalev GI, Buntzman A, Frelinger JA, et al. A Humanized Mouse Model to Study Hepatitis C Virus Infection, Immune Response, and Liver Disease. *Gastroenterology*, in press, doi:10.1053/j.gastro.2011.01.001.

Carine Fillebeen

Lady Davis Institute for Medical Research
Sir Mortimer B. Davis Jewish General Hospital
3755 Cote-Ste-Catherine Road, Montreal
Quebec, Canada H3T 1E2

Kostas Pantopoulos*

Lady Davis Institute for Medical Research
Sir Mortimer B. Davis Jewish General Hospital
3755 Cote-Ste-Catherine Road, Montreal
Quebec, Canada H3T 1E2

Department of Medicine, McGill University, Canada

* Tel.: +1 514 340 8260x5293; fax: +1 514 340 7502

E-mail addresses: kostas.pantopoulos@mcgill.ca